

Turf Supreme® 16-6-8 plus Trimec®

ACTIVE INGREDIENTS:	
2,4-dichlorophenoxyacetic acid	0.55%
(+)-(R)-2-(2 methly-4-chlorophenoxy) propionic acid	0.12%
Dicamba: 3,6-dichloro-o-anisic acid	0.05%
INERT INGREDIENTS:	
TOTAL	100.00%

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

EPA REG. NO. 2217-643-7001

EPA EST. NO.61842-CA-1

THIS PRODUCT CONTAINS:

Isomer Specific by AOAC Methods

11.0 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.

1.0 lbs. 3,6-dichloro-o-anisic acid equivalent per ton.

2.4 lbs. (+)-(R)-2-(2 methyl-4-chlorophenoxy) propionic acid equivalent per ton.

STOP! READ THE ENTIRE LABEL FIRST. **OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY. KEEP OUT OF REACH CHILDREN** CAUTION

PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals

CAUTION: Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing dust. Wear long-sleeved shirt, long pants, shoes, socks and chemical-resistant gloves made of any waterproof material.

F	IRS	Т	Α	D	

 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
 Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
 Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
···· ··· ··· ··

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate. This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

GUARANTEED ANALYSIS: 16-6-8	
TOTAL NITROGEN (N)	
16.0% Ammoniacal Nitrogen	
AVAILABLE PHOSPHATE (P ₂ O ₅)	6.0%
SOLUBLE POTASH (K ₂ O)	
Sulfur (S)	
16.0% Combined Sulfur (S)	
Iron (Fe)	
Manganese (Mn)	
o ()	

Derived from Monoammonium Phosphate, Ammonium Sulfate, Muriate of Potash, Iron Oxysulfate and Manganese Oxysulfate. Information regarding the contents and levels of metals in this product is available on the Internet at http://www.regulatory-info-jr.com

SPREADER SETTINGS: Apply 4 lbs. per 1,000 sq.ft. SGN 150

BEST [®] Pro	4B
Spyker / BEST [®] Models 34, 44, 64	3.7 Rates are for spreading one
Earthway	12 time across in one direction
Lesco	13
Scotts R-8A (Cone 8)	J ¹ / ₂
Lely 1250 / Vicon	visit www.bestfertilizer.com

As a guide, set spreader at the setting indicated by the above chart before applying. Remember the settings provided are only a guide and depend on the age, condition and speed of operation of your spreader. Check spreader calibration often and make adjustment as needed. Keep this product off sidewalks, patios, driveways, etc. This product contains materials which will stain concrete and painted surfaces. If it comes in contact with concrete surface, sweep off immediately before finishing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter product beyond lawn edge and onto concrete. Calibration Instructions. To provide proper distribution, calibrate spreader before application. Calibrate our spreader on 300 sq.ft. and adjust to apply 1.0 lb. of product per 300 sq.ft.

bestfertilizer.com

Simplot Professional Products • P.O. Box 198, Lathrop, CA 95330 • 800-992-6066

Spread The Word

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled. Do not apply this product to dichondra or carpetgrass. Do not apply this product to 'Floratam' St. Augustinegrass in Florida. Do not use this product where desirable clovers are present. Do not exceed specified dosages for any area. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning to prevent over-application. Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result. Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

INSTRUCTIONS

Turf Supreme® 16-6-8 Plus Trimec® controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a 16-6-8 fertilizer provides plant nutrients or a supply of plant food. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days.

The maximum broadcast application rate is 5.0 pounds of product per 1,000 sq. ft. per application (1.2 lb. 2,4-D ae, 0.27 lb MCPP-p ae and 0.11 lb. dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 10.0 pounds of product per 1,000 sq.ft. (2.4 lb. 2,4-D ae, 0.54 lb. MCPP-p ae and 0.22 lb. dicamba ae per acre).

- 1. Mow the lawn to normal height 1 to 2 days before application.
- 2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
- Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
- 4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

WEEDS CONTROLLED

Annual yellow sweetclover Bindweed, field Bird vetch Bitter wintercress Bittercress, hairy Black medic Black mustard Blackseed plantain Blessed thistle Bloodflower milkweed Bracted plantain Bristly oxtongue Broadleaf dock	Broadleaf plantain Buckhorn plantain Bulbous buttercup Bull thistle Burdock Burning nettle Burclover Buttercup Buttercup Buttonweed Canada thistle Carpetweed Chickweed, common	Chickweed, mouseear Chicory Clover, crimson Clover, white Cocklebur Common mullein Curly dock Daisy, English Dandelion Dichondra Dollarweed False dandelion Fiddleneck	Goldenrod Ground ivy Henbit Hoary cress Hoary plantain Kochia Knotweed Lambsquarters Mallow Morningglory Mustard Narrowleaf plantain Narrowleaf vetch	Nettle Oxalis Pearlwort Pigweed Plantain Poison ivy Poison oak Prickly lettuce Prostrate pigweed Prostrate knotweed Prostrate spurge Purslane common Puncturevine	Pigweed Red sorrel Redroot pigweed Redstem filaree Rough cinquefoil Rough fleaban Russian pigweed Russian thistle Scarlet pimpernel Scotch thistle Shepherdspurse Smooth dock Smooth pigweed	Sowthistle Speedwell Spotted spurge Stinging nettle Thistle Venice mallow White mustard Wild carrot Yarrow

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTIAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

PACKAGE SIZE:

50-LB. (22.68-Kg.) BAGS MANUFACTURED IN U.S.A. 74070 (PRO/TB R9-14-2010)

HANDLING PRECAUTIONS: MSDS - See Reverse CAUTION: KEEP OUT OF THE REACH OF CHILDREN. HARMFUL IF SWALLOWED - DO NOT INHALE